

Patient information from BMJ

Last published: Sep 17, 2020

Diarrhoea in adults

Diarrhoea often clears up on its own after a few days. But if you feel very ill, you may need to see a doctor.

This information looks at diarrhoea in adults. Some of the treatments we talk about aren't recommended for children.

What is diarrhoea?

If you have diarrhoea you have loose, watery stools or you need to go to the toilet a lot more often than normal.

This information looks at diarrhoea that's caused by an infection with a virus or bacteria. You can pick these up from contaminated food or drink.

You can also catch an infection that can cause diarrhoea from another person who has it. This happens when you touch something that's contaminated with traces of their stool (for example, the person's hand) and then put your hand to your mouth (for example, when you eat).

People often get diarrhoea when they're travelling in other countries (called travellers' diarrhoea). It's a good idea to pack diarrhoea medicine whenever you go on holiday.

What are the symptoms?

As well as watery stools, you may also get vomiting or stomach cramps. Because you lose water in your stools, having diarrhoea means your body may lose too much fluid (you get dehydrated). This makes you feel thirsty, and you might have dark-coloured urine, or feel sleepy.

If your diarrhoea doesn't clear up in a few days you should see a doctor. You should also see a doctor if:

- you have a high temperature (fever)
- there's blood in your stools
- you've been travelling abroad

Diarrhoea in adults

- you've just come out of hospital
- you work as a carer, or
- you work in a job where you handle food.

If you're too ill to eat or drink, see a doctor as soon as you can.

What treatments work?

Prevention

You can help prevent diarrhoea by washing your hands after using the toilet, before preparing food, and before eating.

There are other precautions you can take when travelling abroad, especially in countries where tap water might not be safe to drink.

When travelling in these countries you should only drink water that is bottled and properly sealed. Avoid eating or drinking:

- ice and tap water
- salads, previously peeled fruits, and raw foods
- unpackaged condiments and sauces
- food from street vendors and buffets.

Taking care of yourself

Some people prefer to eat simple foods, like bread, if they have diarrhoea. But the most important thing is to have **plenty to drink**.

It's best to avoid drinks that contain a lot of sugar as they can make diarrhoea worse. You should also avoid drinks that contain caffeine.

You can also go to a pharmacy and buy **rehydration salts** that you mix with water. They are designed to replace fluids and mineral salts your body may have lost. But, for people with mild dehydration, it's not clear whether they're better for rehydration than water or soft drinks.

Medicines

There are two main types of medicine that can help if you have diarrhoea: antibiotics to fight bacterial infection, and drugs that slow down your bowel movements.

Antibiotics are drugs that kill bacteria. But not all diarrhoea is caused by bacteria, so antibiotics aren't the right treatment for everyone. Your doctor may do tests to see what kind of diarrhoea you have.

Antibiotics also have side effects. Some of these side effects may be similar to the symptoms you want treated, such as nausea, a stomach ache, or diarrhoea. But they aren't usually serious enough for people to stop taking their antibiotics.

Diarrhoea in adults

Some medicines help to deal with diarrhoea by slowing down your bowel movements. You can buy some of these from a pharmacy.

But they're not recommended for severe diarrhoea, as they can mask your symptoms. This means that, because they reduce your diarrhoea, you seem more recovered than you are.

These medicines are most useful in people whose symptoms are not severe, or who have begun to recover and who possibly need to travel.

Some of these medicines can cause side effects, including dizziness and constipation.

You shouldn't take these medicines if you have a fever or if there's blood or mucus in your stools. If you have these symptoms, see a doctor.

What will happen to me?

Diarrhoea isn't usually serious. Most people get better on their own without any problems.

The main risk is losing too much water (getting dehydrated). So it's important to drink plenty of fluids, especially if you're an older person. Drinking often, in small sips, may be the best way to drink with an upset stomach. Occasionally, people who get very dehydrated need to go to hospital.

The patient information from *BMJ Best Practice* from which this leaflet is derived is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2020. All rights reserved.

